



NUMERICAL INVESTIGATION OF WAVE EQUATIONS IN LARGE DOMAINS VIA A NOVEL VARIATIONAL ITERATION METHOD

HOSSEIN GHANEAI*, MOHAMMAD MIRABI, AND REZA RASHIDI MEYBODI

ABSTRACT. The value of an auxiliary parameter incorporated into the well-known variational iteration method (VIM) to obtain solutions of wave equations in unbounded domains is discussed in this article. The suggested method, namely the optimal variational iteration method, is investigated for convergence. Furthermore, the proposed method is tested on one-dimensional and two-dimensional wave equations in unbounded domains in order to better understand the solution mechanism and choose the best auxiliary parameter. Comparisons with results from the standard variational iteration procedure demonstrate that the auxiliary parameter is very useful in tracking the convergence field of the approximate solution.

Keywords: Wave equations, Unbounded domains, Variational iteration method, Optimal variational iteration method, Auxiliary parameter, Hermite-Gauss quadrature.

1. Introduction

Nonlinear partial differential equations can be used to describe a wide range of phenomena in a variety of fields. Most of these equations do not have a precise analytical solution, so these nonlinear equations should be solved by approximate methods. Since most of these equations do not have a direct empirical solution, they must be solved using approximate methods. Wave equations are the most important of these equations, so we concentrated our research on them. Waves appear in unbounded media in a variety of areas, including vibrations, aerodynamics, solid geophysics, oceanography, meteorology, and electromagnetics. In unbounded domains, the wave equation is as follows:

$$(1.1) \quad \frac{\partial^2 \mathbf{u}(\mathbf{x}, t)}{\partial t^2} - c^2 \Delta \mathbf{u}(\mathbf{x}, t) = 0,$$

subject to the initial conditions

$$(1.2) \quad \begin{cases} \mathbf{u}(\mathbf{x}, 0) = f(\mathbf{x}) \\ \mathbf{u}_t(\mathbf{x}, 0) = g(\mathbf{x}), \end{cases} \quad \mathbf{x} \in \mathbb{R}^n.$$

Since the 1970s, numerical approaches for such problems have been developed[15]. The key categories of methods that have appeared are boundary integral methods, infinite element methods, absorbing layer methods, and non-reflecting boundary state (NRBC) methods [13, 23, 24, 25, 26, 27, 28, 29]. The standard method for solving such problems numerically

Date: Received: December 25, 2021, Accepted: May 7, 2022.

*Corresponding author.

is to create an imaginary boundary and add acceptable boundary conditions [2, 14, 20, 19]. The process of integral equations is also a useful technique for converting this partial differential equation to an integral equation on the scatterer's bounded surface [9, 34]. Laplace-Fourier methods with Galerkin boundary elements in space and collocation methods with some stabilization techniques are some of the latest numerical methods [3, 5, 6, 7, 17, 18]. In Ref. [8] a quick version of the marching-on-in-time (MOT) approach is proposed, which is based on a suitable plane wave expansion of the arising potential, resulting in lower computational and storage costs [1]. Wazwaz used the Adomian decomposition approach to present approximate analytical solutions to wave equations in the unbounded domain. Furthermore, the Homotopy perturbation technique is used to approximate wave equation solutions in an unbounded domain [40, 4].

The variational iteration method is a more efficient and convenient analytical procedure. In Ref. [21], Ji-Huan He is a Chinese mathematician. He presented a very clear and basic explanation of the variational iteration process, which was further developed by the inventor himself. [39, 22, 37]. The method's key feature is its adaptability and ability to obtain reliable and convenient solutions to nonlinear equations [38, 16, 33, 10]. Furthermore, there are many variations of the variational iteration process, the most appealing of which is Herisanu and Marinca's, in which the variational iteration method is combined with least squares technology, and one iteration leads to perfect outcomes [22]. Yilmaz and Inc developed a variational iteration algorithm with an auxiliary parameter to change the convergence rate, but they did not have a general rule for the right auxiliary parameter selection [10]. Hosseini et al. improved this updated approach by adding several profitable rules for determining the auxiliary parameter optimally [11, 35, 32, 12].

The variational iteration approach with an auxiliary parameter, optimum variational, is successfully used in the present paper to achieve approximate solutions of wave equations in unbounded domains. The residual function and the residual function error are described in the proposed method to select the auxiliary parameter optimally.

2. The variational iteration method

The basic solution protocol of the variational iteration process is briefly recapitulated here. Consider the following equation in terms of function:

$$(2.1) \quad Hu = Lu + Nu + Ru + g(x),$$

where L represents the higher order derivative, which is thought to be conveniently invertible. R stands for a linear differential operator with a lower order than L , Nu for nonlinear terms, and g for the root name. He's approach is characterized by the construction of a correction functional for the equation (2.1), which reads as follows:

$$(2.2) \quad u_{n+1}(x) = u_n(x) + \int_0^x \lambda(s) Hu_n(s) ds,$$

where λ is a Lagrange multiplier which can be identified optimally via variational theory [39], u_n is the n th approximate solution, and \tilde{u}_n denotes a restricted variation, i.e., $\partial\tilde{u}_n = 0$. After identification of the multiplier, a variational iteration algorithm is constructed, an exact solution can be achieved when n tends to infinite:

$$(2.3) \quad u(x) = \lim_{n \rightarrow \infty} u_n(x).$$

In summary, we have the following variational iteration formula for **(2.1)**:

$$(2.4) \quad \begin{cases} u_0(x) \text{ is an arbitrary function,} \\ u_{n+1}(x) = u_n(x) + \int_0^x \lambda(s) H u_n(s) ds, \end{cases} \quad n \geq 0.$$

3. Optimal Variational Iteration Method

In the variational iteration procedure, equation **(2.4)**, an unknown auxiliary parameter can be included:

$$(3.1) \quad \begin{cases} u_0(x) \text{ is an arbitrary function,} \\ u_1(x, h) = u_0(x) + h \int_0^x \lambda(s) H u_n(s) ds, \\ u_{n+1}(x, h) = u_n(x, h) + h \int_0^x \lambda(s) H u_n(s, h) ds, \end{cases} \quad n \geq 1.$$

It should be noted that symbolic computation tools like Maple or Mathematica can compute $u_{n+1}(x, h)$, $n \geq 1$. The auxiliary parameter h is included in the approximate solutions $u_{n+1}(x, h)$, $n \geq 1$. The method's validity is based on the premise that the approximation $u_{n+1}(x, h)$, $n \geq 1$, converges to the exact solution. It is the auxiliary parameter that guarantees the assumption is met. In general, using the residual function's error of norm two, it is simple to select an appropriate value of h that assures the approximation solutions are convergent [10, 11, 35, 32]. In reality, the suggested technique is relatively basic, easier to implement, and capable of more correctly approximating the solution in a broad solution domain.

4. Convergence analysis

In this section, the convergence of the optimal variational iteration method is studied according to the alternative approach of this method which present in the following.

This approach can be implemented, in a reliable and efficient way, to handle the nonlinear differential equation **(2.1)**. The linear operator L is defined as $L = \frac{\partial^2}{\partial t^2}$, when the optimal variational iteration method is applied to solve the wave equation **(1.1)**.

Now, define the operator A as,

$$(4.1) \quad A u(\mathbf{x}, t, h) = h \int_0^t \lambda(\tau) H u(\mathbf{x}, \tau, h) d\tau,$$

and define the components v_n, s_n , $n \geq 0$, as,

$$\begin{cases} v_0(\mathbf{x}, t) = u_0(\mathbf{x}, t), \\ s_0(\mathbf{x}, t) = v_0(\mathbf{x}, t), \\ \\ v_1(\mathbf{x}, t, h) = A s_0(\mathbf{x}, t), \\ s_1(\mathbf{x}, t, h) = s_0(\mathbf{x}, t) + v_1(\mathbf{x}, t, h), \end{cases}$$

and in general for $n \geq 1$,

$$(4.2) \quad \begin{cases} v_{n+1}(\mathbf{x}, t, h) = A s_n(\mathbf{x}, t, h), \\ s_{n+1}(\mathbf{x}, t, h) = s_n(\mathbf{x}, t, h) + v_{n+1}(\mathbf{x}, t, h), \end{cases}$$

then, consequently, we have,

$$(4.3) \quad u(\mathbf{x}, t, h) = \lim_{n \rightarrow \infty} s_n(\mathbf{x}, t, h) = v_0(\mathbf{x}, t) + \sum_{n=1}^{\infty} v_n(\mathbf{x}, t, h).$$

The zeroth approximation $u_0(\mathbf{x}, t)$ can be freely chosen if it satisfies the initial conditions of the problem and $Lu_0(\mathbf{x}, t) = 0$. For the approximation purpose, we approximate the solution $u(\mathbf{x}, t, h) = v_0(\mathbf{x}, t) + \sum_{n=1}^{\infty} v_n(\mathbf{x}, t, h)$, by the N th-order truncated series $u_N(\mathbf{x}, t, h) = v_0(\mathbf{x}, t) + \sum_{n=1}^N v_n(\mathbf{x}, t, h)$.

The approximate solutions $u_N(t, h)$, contains the auxiliary parameter h . It is the auxiliary parameter that ensures that the assumption can be satisfied, in general, by means of the error of norm two of the residual function.

The sufficient conditions for convergence of the method and the error estimate will be introduced. The main results are proposed in the following theorems [36].

Theorem 4.1 Let A , defined in (4.1), be an operator from a Hilbert space H to H . If $\exists \tilde{h} \neq 0, 0 < \gamma < 1$, such that,

$$\begin{cases} \|As_0(\mathbf{x}, t)\| \leq \gamma \|s_0(\mathbf{x}, t)\|, \\ \|As_1(\mathbf{x}, t, \tilde{h})\| \leq \gamma \|As_0(\mathbf{x}, t)\|, \\ \|As_n(\mathbf{x}, t, \tilde{h})\| \leq \gamma \|As_{n-1}(\mathbf{x}, t, \tilde{h})\|, \end{cases} \quad n = 2, 3, 4, \dots$$

Then the series solution defined in (4.3),

$$u(\mathbf{x}, t) = \lim_{n \rightarrow \infty} s_n(\mathbf{x}, t, \tilde{h}) = v_0(\mathbf{x}, t) + \sum_{n=1}^{\infty} v_n(\mathbf{x}, t, \tilde{h}),$$

converges [11].

Lemma 4.1 Let L , defined in (2.1), be as follow as, $L = \frac{\partial^2}{\partial t^2}$, and λ identified optimally via variational theory [11]. If k , be a function from a Hilbert space H to H , then we have,

$$L \left\{ \int_0^t \lambda(\tau) k(\mathbf{x}, \tau) d\tau \right\} = -k(\mathbf{x}, t).$$

Theorem 4.2 Let L , defined in (2.1), be as follow as, $L = \frac{\partial^2}{\partial t^2}$, if we have $u(\mathbf{x}, t) = v_0(\mathbf{x}, t) + \sum_{n=1}^{\infty} v_n(\mathbf{x}, t, \tilde{h})$, then $u(\mathbf{x}, t)$, is an exact solution of the nonlinear problem (2.1).

Theorem 4.3 Suppose that the series solution $u(\mathbf{x}, t) = v_0(\mathbf{x}, t) + \sum_{n=1}^{\infty} v_n(\mathbf{x}, t, \tilde{h})$, defined in (4.3), is convergent to exact solution of the nonlinear problem (2.1). If the truncated series $u_N(\mathbf{x}, t) = v_0(\mathbf{x}, t) + \sum_{n=1}^N v_n(\mathbf{x}, t, \tilde{h})$, is used as an approximate solution, then the maximum error is estimated as,

$$\|u(\mathbf{x}, t) - u_N(\mathbf{x}, t)\| \leq \frac{1}{1 - \gamma} \gamma^{N+1} \|v_0\|.$$

In summary, we can define,

$$\beta_i = \begin{cases} \frac{\|v_{i+1}\|}{\|v_i\|}, & \|v_i\| \neq 0, \\ 0, & \|v_i\| = 0, \end{cases} \quad i = 0, 1, 2, \dots$$

Now, if $0 < \beta_i < 1$ for $i = 0, 1, 2, \dots$, then the series solution $v_0(\mathbf{x}, t) + \sum_{n=1}^{\infty} v_n(\mathbf{x}, t, \tilde{h})$, of problem (2.1) converges to an exact solution, $u(\mathbf{x}, t)$. Moreover, as stated in Theorem 4.3,

the maximum absolute truncation error is estimated to be,

$$\| u(\mathbf{x}, t) - u_N(\mathbf{x}, t) \| \leq \frac{1}{1 - \beta} \beta^{N+1} \| v_0 \|,$$

where $\beta = \max \{ \beta_i, i = 0, 1, 2, \dots \}$.

Notice that, the first finite terms do not affect the convergence of series solution. In other words, if the first finite β_i 's, $i = 0, 1, 2, \dots, l$, are not less than one and $\beta_i < 1$, for $i > l$, then, of course the series solution $v_0(\mathbf{x}, t) + \sum_{n=1}^{\infty} v_n(\mathbf{x}, t, \tilde{h})$, of problem (2.1), converges to an exact solution [36].

5. Numerical Examples

To elucidate the solution procedure, three examples are given. In each case, the standard variational iteration method is applied on the wave equations in unbounded domains. The obtained results show that where the solutions of wave equations in unbounded domains are under investigation, the standard variational iteration method is not applicable. Therefore, the proposed method is tested on the aforementioned wave equation in unbounded domains. Comparison with results by exact solutions indicates that the large domains will not decrease the effectiveness of the proposed method.

Example 5.1 Consider the following wave equation [38]:

$$(5.1) \quad \begin{cases} u_{tt} = u_{xx}, & -\infty < x < \infty, & t > 0, \\ u(x, 0) = \sin(x), & u_t(x, 0) = 0, & -\infty < x < \infty, \end{cases}$$

which admits the solution $u(x, t) = \sin(x) \cos(t)$. Take $(x, t) \in (-\infty, \infty) \times [0, 50]$. According to the standard VIM we have the following variational iteration formula:

$$(5.2) \quad u_{n+1}(x, t) = u_n(x, t) + \int_0^t (s - t) \left\{ \frac{\partial^2 u_n(x, s)}{\partial s^2} - \frac{\partial^2 u_n(x, s)}{\partial x^2} \right\} ds.$$

Beginning with $u_0(x, t) = u(x, 0) + tu_t(x, 0) = \sin(x)$, we stop the solution procedure at $u_{60}(x, t)$. Figure 1, is the absolute error of $u_{60}(x, t)$, for $(x, t) \in [-10^8, 10^8] \times [0, 50]$, showing that the solution $u_{60}(x, t)$ is not valid for large values of x and t , of course, the accuracy can be improved if the iteration procedure continues and the exact solution can be obtained when n tends to infinite. Now, using the recursive scheme (3.1), we successively have:

$$\begin{aligned} u_0(x, t) &= u(x, 0) + tu_t(x, 0) = \sin(x), \\ u_1(x, t) &= \sin(x) - \frac{1}{2}ht^2 \sin(x), \end{aligned}$$

and in general,

$$(5.3) \quad \begin{aligned} u_{n+1}(x, t, h) &= u_n(x, t, h) \\ &+ h \int_0^t (s - t) \left\{ \frac{\partial^2 u_n(x, s, h)}{\partial s^2} - \frac{\partial^2 u_n(x, s, h)}{\partial x^2} \right\} ds, \quad n \geq 1. \end{aligned}$$

In order to find a proper value of h for the approximate solutions (5.3), we define the following residual function,

$$(5.4) \quad r_{60}(x, t, h) = \frac{\partial^2 u_{60}(x, t, h)}{\partial t^2} - \frac{\partial^2 u_{60}(x, t, h)}{\partial x^2},$$

and the following error of residual function,

$$(5.5) \quad e_{60}(h) = \int_{-\infty}^{\infty} \int_0^{50} |r_{60}(x, t, h)|^2 dt dx.$$

We apply Hermite-Gauss quadrature as a numerical integration to calculate the approximate $e_{60}(h)$ [31]. For obtaining an optimal value of h , we choose the minimum point of the error residual function (5.4). The minimum point of $e_{60}(h)$, as $h = 0.95077$, is obtained by using optimization package of Maple software in 2 seconds. By substituting $h = 0.95077$, in $u_{60}(x, t, h)$, the absolute error of the 60th-order approximation of the proposed method reduces remarkably, as shown in Figure 2.




FIGURE 1. Absolute error for the 60th-order approximation by standard VIM for $u(x, t)$ in example 5.1.




FIGURE 2. Absolute error for the 60th-order approximation by optimal VIM when $h = 0.95077$, in example 5.1.

Example 5.2 Consider the following wave equation [38]:

$$(5.6) \quad \begin{cases} u_{tt} = u_{xx}, & -\infty < x < \infty, & t > 0. \\ u(x, 0) = \sin(x), & u_t(x, 0) = \cos(x), & -\infty < x < \infty, \end{cases}$$

It is easy to verify that $u(x, t) = \sin(x+t)$. We take the solution domain as $(x, t) \in (-\infty, \infty) \times [0, 100]$. Similarly the absolute error of $u_{120}(x, t)$ for $(x, t) \in [-10^8, 10^8] \times [0, 100]$, tends to very large values, when time tends to 100, as illustrated in Figure 3. Using the iteration formulation (3.1), we successively have

$$\begin{aligned} u_0(x, t) &= u(x, 0) + tu_t(x, 0) = \sin(x) + t \cos(x), \\ u_1(x, t) &= \sin(x) + t \cos(x) + \frac{1}{6}h(t^3 \cos(x) + 3t^2 \sin(x)), \end{aligned}$$

and in general,

$$(5.7) \quad \begin{aligned} u_{n+1}(x, t, h) &= u_n(x, t, h) \\ &+ h \int_0^t (s-t) \left\{ \frac{\partial^2 u_n(x, s, h)}{\partial s^2} - \frac{\partial^2 u_n(x, s, h)}{\partial x^2} \right\} ds, \quad n \geq 1. \end{aligned}$$

We define the residual function of $u_{120}(x, t)$ as,

$$(5.8) \quad r_{120}(x, t, h) = \frac{\partial^2 u_{120}(x, t, h)}{\partial t^2} - \frac{\partial^2 u_{120}(x, t, h)}{\partial x^2}.$$

For obtaining an optimal value of h , we choose the global minimum point of the error of residual function (5.8):

$$(5.9) \quad e_{120}(h) = \int_{-\infty}^{\infty} \int_0^{100} |r_{120}(x, t, h)|^2 dt dx.$$

Thus, we select $h = 0.93258$, and it's absolute error is reduced greatly for $u_{120}(x, t)$, $(x, t) \in [-10^8, 10^8] \times [0, 100]$, as shown in Figure 4.




FIGURE 3. Absolute error for the 120th-order approximation by standard VIM for $u(x, t)$ in example 5.2.




FIGURE 4. Absolute error for the 120th-order approximation by optimal VIM when $h = 0.93258$, in example 5.2.

Example 5.3 Consider the following two-dimensional wave equation [4]:

$$(5.10) \quad \begin{cases} u_{tt} = 2(u_{xx} + u_{yy}), & -\infty < x, y < \infty, \quad t > 0, \\ u(x, y, 0) = \sin(x) \sin(y), \quad u_t(x, y, 0) = 0, & -\infty < x, y < \infty, \end{cases}$$

which admits the solution $u(x, y, t) = \sin(x) \sin(y) \cos(2t)$. Take $(x, y, t) \in (-\infty, \infty) \times (-\infty, \infty) \times [0, 50]$ According to the standard VIM we have the following variational iteration formula:

$$(5.11) \quad \begin{aligned} u_{n+1}(x, y, t) &= u_n(x, y, t) \\ &+ \int_0^t (s-t) \left\{ \frac{\partial^2 u_n(x, y, s)}{\partial s^2} - 2 \left(\frac{\partial^2 u_n(x, y, s)}{\partial x^2} + \frac{\partial^2 u_n(x, y, s)}{\partial y^2} \right) \right\} ds. \end{aligned}$$

Beginning with $u_0(x, y, t) = u(x, y, 0) + tu_t(x, y, 0) = \sin(x) \sin(y)$, we stop the solution procedure at $u_{120}(x, y, t)$. Figure 5, is the absolute error of $u_{120}(x, 10^8, t)$, for $(x, t) \in [-10^8, 10^8] \times [0, 50]$, showing that the solution $u_{120}(x, y, t)$ is not valid for large values of x, y and t , of course, the accuracy can be improved if the iteration procedure continues and the exact solution can be obtained when n tends to infinite. Now, using the recursive scheme (3.1), we successively have:

$$\begin{aligned} u_0(x, y, t) &= u(x, y, 0) + tu_t(x, y, 0) = \sin(x) \sin(y), \\ u_1(x, y, t, h) &= \sin(x) \sin(y) - ht^2 \sin(x) \sin(y), \end{aligned}$$

and in general,

$$(5.12) \quad u_{n+1}(x, y, t, h) = u_n(x, y, t) + h \int_0^t (s-t) \left\{ \frac{\partial^2 u_n(x, y, s)}{\partial s^2} - 2 \left(\frac{\partial^2 u_n(x, y, s)}{\partial x^2} + \frac{\partial^2 u_n(x, y, s)}{\partial y^2} \right) \right\} ds, \quad n \geq 1.$$

In order to find a proper value of h for the approximate solutions (5.12), we define the following residual function,

$$r_{120}(x, y, t, h) = \frac{\partial^2 u_{120}(x, y, t, h)}{\partial t^2} - 2 \left(\frac{\partial^2 u_{120}(x, y, t, h)}{\partial x^2} + \frac{\partial^2 u_{120}(x, y, t, h)}{\partial y^2} \right),$$

and the following error of residual function,

$$e_{120}(h) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_0^{50} |r_{120}(x, y, t, h)|^2 dt dy dx,$$

clearly, suitable value of h , is the global minimum point of $e_{120}(h)$ which we obtained $h = 0.93253$, using Maple software in 5 seconds.. The absolute error of 120th-order approximation of the proposed method for $u_{120}(x, 10^8, t)$ in the solution domain $(x, t) \in [-10^8, 10^8] \times [0, 50]$, is given in Figure 6, the accuracy is remarkably improved by the optimal choice of h .




FIGURE 5. Absolute error for the 120th-order approximation by standard VIM for $u_{120}(x, 10^8, t)$ in example 5.3.




FIGURE 6. Absolute error for the 120th-order approximation by present technique for $u_{120}(x, 10^8, t)$ when $h = 0.93253$ in example 5.3.

Conclusion

Many application problems have been successfully solved using the variational iteration process. The focus of this article is on how the propounded approach may manage solutions of wave equations in large domains, while the standard VIM may have difficulty obtaining adequate precision in large domains. It is demonstrated that VIM with an auxiliary parameter is a powerful and efficient tool for wave equations. Comparing the original VIM, our approach is simple to apply and capable of approximating solutions more precisely over a longer interval. In particular, the proposed approach widens the convergence zone. Furthermore, the

proposed approach can be conveniently extended to a broad range of nonlinear problems in large domains.

REFERENCES

- [1] Banjai, L., and Stefan Sauter. Rapid solution of the wave equation in unbounded domains. *SIAM Journal on Numerical Analysis* 47.1 (2008): 227-249.
- [2] Bayliss, Alvin, and Eli Turkel. "Radiation boundary conditions for wave-like equations." *Communications on Pure and applied Mathematics* 33, no. 6 (1980): 707-725.
- [3] Birgisson, B., E. Siebrits, and A. P. Peirce. Elastodynamic direct boundary element methods with enhanced numerical stability properties. *International journal for numerical methods in engineering* 46.6 (1999): 871-888.
- [4] Chun, Changbum, Hossein Jafari, and Yong-II Kim. Numerical method for the wave and nonlinear diffusion equations with the homotopy perturbation method. *Computers and Mathematics with Applications* 57.7 (2009): 1226-1231.
- [5] Davies, Penny J. Numerical stability and convergence of approximations of retarded potential integral equations. *SIAM journal on numerical analysis* 31.3 (1994): 856-875.
- [6] Davies, Penny J., and Dugald B. Duncan. Stability and convergence of collocation schemes for retarded potential integral equations. *SIAM journal on numerical analysis* 42.3 (2004): 1167-1188.
- [7] Ding Y, Forestier A, Duong TH. A Galerkin scheme for the time domain integral equation of acoustic scattering from a hard surface. In *The Journal of the Acoustical Society of America CONF 4320FN 1989 Oct* (Vol. 86, Nu. 4, pp. 1566-1572). Acoustical Society of America.
- [8] Ergin, A. A., B. Shanker, and E. Michielssen. Fast analysis of transient acoustic wave scattering from rigid bodies using the multilevel plane wave time domain algorithm. *The Journal of the Acoustical Society of America* 107 (2000): 1168-1178.
- [9] Friedman, M. B., and R. Shaw. Diffraction of pulses by cylindrical obstacles of arbitrary cross section. *Journal of Applied Mechanics* 29 (1962): 40.
- [10] Ghaneai, H., and M. M. Hosseini. "Solving differential-algebraic equations through variational iteration method with an auxiliary parameter." *Applied Mathematical Modelling* 40, no. 5-6 (2016): 3991-4001.
- [11] Ghaneai, H., and M. M. Hosseini. "Variational iteration method with an auxiliary parameter for solving wave-like and heat-like equations in large domains." *Computers & Mathematics with Applications* 69, no. 5 (2015): 363-373.
- [12] H. Ghaneai, M.M. Hosseini, S.T. Mohyud-Din, Modified variational iteration method for solving a neutral functional-differential equation with proportional delays, *Int. J. Numer. Method H.* 22.8 (2012) 1086-1095.
- [13] Givoli, Dan. "High-order local non-reflecting boundary conditions: a review." *Wave motion* 39, no. 4 (2004): 319-326.
- [14] Ghaneai, H., M. M. Hosseini, and Syed Tauseef Mohyud-Din. "Modified variational iteration method for solving a neutral functional-differential equation with proportional delays." *International Journal of Numerical Methods for Heat & Fluid Flow* (2012).
- [15] Givoli, Dan. *Numerical methods for problems in infinite domains*. Elsevier, 2013.
- [16] Gonzalez-Gaxiola, O., Anjan Biswas, Mehmet Ekici, and Salam Khan. "Highly dispersive optical solitons with quadratic-cubic law of refractive index by the variational iteration method." *Journal of Optics* (2021): 1-8.
- [17] M Abolhasani, H Ghaneai, M Heydari. Modified homotopy perturbation method for solving delay differential equations. *Appl. Sci. Reports*, 16.2 (2010): 89-92.
- [18] Hosseini, M. M., et al. "Tri-prong scheme for regularized long wave equation." *Journal of the Association of Arab Universities for Basic and Applied Sciences* 20 (2016): 68-77.
- [19] Hosseini, M. M., et al. "Auxiliary parameter in the variational iteration method and its optimal determination." *International Journal of Nonlinear Sciences and Numerical Simulation* 11.7 (2010): 495-502.
- [20] Hosseini, Said Mohammad Mehdi, Syed Tauseef Mohyud-Din, and Husain Ghaneai. "Variational iteration method for Hirota-Satsuma coupled KdV equation using auxiliary parameter." *International Journal of Numerical Methods for Heat & Fluid Flow* (2012).
- [21] J.H. He, Variational iteration method—a kind of non-linear analytical technique: some examples, *Int. J. Nonlinear Mech.* 34.4 (1999): 699-708.

- [22] J.H. He, Variational iteration method-Some recent results and new interpretations, *J. Comput. Appl. Math.* 207 (2007) 3-17.
- [23] Heydari, M., et al. "A novel hybrid spectral-variational iteration method (HS-VIM) for solving nonlinear equations arising in heat transfer." (2013): 501-512.
- [24] Heydari, M., et al. "Solution of strongly nonlinear oscillators using modified variational iteration method." *International Journal of Nonlinear Dynamics in Engineering and Sciences* 3 (2011): 33-45.
- [25] Heydari, Mohammad, Ghasem Barid Loghmani, and Abdul-Majid Wazwaz. "A numerical approach for a class of astrophysics equations using piecewise spectral-variational iteration method." *International Journal of Numerical Methods for Heat & Fluid Flow* (2017).
- [26] Nikooeinejad, Z., and Mohammad Heydari. "Nash equilibrium approximation of some class of stochastic differential games: A combined Chebyshev spectral collocation method with policy iteration." *Journal of Computational and Applied Mathematics* 362 (2019): 41-54.
- [27] Nikooeinejad, Z., Ali Delavarkhalafi, and Mohammad Heydari. "Application of shifted Jacobi pseudospectral method for solving (in) finite-horizon min-max optimal control problems with uncertainty." *International Journal of Control* 91.3 (2018): 725-739.
- [28] Avazzadeh, Z., et al. "Smooth solution of partial integro-differential equations using radial basis functions." *The Journal of Applied Analysis and Computation* 4.2 (2014): 115-127.
- [29] Hosseini, M. M., et al. "Study on hyperbolic telegraph equations by using homotopy analysis method." *Studies in Nonlinear Sciences* 1.2 (2010): 50-56.
- [30] Rehman, Shahida, Akhtar Hussain, Jamshaid Ul Rahman, Naveed Anjum, and Taj Munir. "Modified Laplace based variational iteration method for the mechanical vibrations and its applications." *acta mechanica et automatica* 16, no. 2 (2022).
- [31] J. Shen, T. Tang, L.L. Wang, *Spectral Methods: Algorithms, Analysis and Applications*. Springer, New York, 2011.
- [32] M.M. Hosseini, S.T. Mohyud-Din, H. Ghaneai, On the coupling of auxiliary parameter, Adomian's polynomials and correction functional, *Math. Comput. Appl.* 16.4 (2011) 959.
- [33] Nikooeinejad, Z., M. Heydari, and G. B. Loghmani. "Numerical solution of two-point BVPs in infinite-horizon optimal control theory: a combined quasilinearization method with exponential Bernstein functions." *International Journal of Computer Mathematics* 98, no. 11 (2021): 2156-2174.
- [34] Sauter, Stefan, and Christoph Schwab. *Randelementmethoden: Analyse, Numerik und Implementierung schneller Algorithmen*. Springer-Verlag, 2004.
- [35] S.M.M. Hosseini, S.T. Mohyud-Din, H. Ghaneai, Variational iteration method for nonlinear Age-Structured population models using auxiliary parameter, *Z. Naturforsch A* 65.12 (2010) 1137.
- [36] Odibat, Z. M. (2010). A study on the convergence of variational iteration method. *Mathematical and Computer Modelling*, 51(9), 1181-1192.
- [37] Rehman, Shahida, Akhtar Hussain, Jamshaid Ul Rahman, Naveed Anjum, and Taj Munir. "Modified Laplace based variational iteration method for the mechanical vibrations and its applications." *acta mechanica et automatica* 16, no. 2 (2022).
- [38] Reddy, G. Janardhana, Ashwini Hiremath, Mahesh Kumar, O. Anwar Bég, and Ali Kadir. "Unsteady magnetohydrodynamic couple stress fluid flow from a shrinking porous sheet: Variational iteration method study." *Heat Transfer* 51, no. 2 (2022): 2219-2236.
- [39] Rehman, Gohar, Shengwu Qin, Qura Tul Ain, Zaheen Ullah, Muhammad Zaheer, Muhammad Afnan Talib, Qaiser Mehmood, and Muhammad Yousuf Jat Baloch. "A study of moisture content in unsaturated porous medium by using homotopy perturbation method (HPM) and variational iteration method (VIM)." *GEM-International Journal on Geomathematics* 13, no. 1 (2022): 1-10.
- [40] Wazwaz, Abdul-Majid. A reliable technique for solving the wave equation in an infinite one-dimensional medium. *Applied Mathematics and Computation* 92.1 (1998): 1-7.

(Hossein Ghaneai) DEPARTMENT OF COMPUTER ENGINEERING, MEYBOD UNIVERSITY, MEYBOD, IRAN.

Email address: h.ghaneai@meybod.ac.ir

(Mohammad Mirabi) DEPARTMENT OF INDUSTRIAL ENGINEERING, MEYBOD UNIVERSITY, MEYBOD, IRAN.

Email address: mirabi@meybod.ac.ir

(Reza Rashidi Meybodi) DEPARTMENT OF COMPUTER ENGINEERING, MEYBOD UNIVERSITY, MEYBOD, IRAN.

Email address: rashidi@meybod.ac.ir