



SOME REMARKS ON THE PAPER "GLOBAL OPTIMIZATION IN METRIC SPACES WITH PARTIAL ORDERS"

MOOSA GABELEH* AND JACK MARKIN

ABSTRACT. The aim of this note is to show that the main conclusion of a recent paper by Sadiq Basha [S. Sadiq Basha, Global optimization in metric spaces with partial orders, *Optimization*, 63 (2014), 817-825] can be obtained as a consequence of corresponding existing results in fixed point theory in the setting of partially ordered metric spaces. Moreover, by a similar approach, we prove that in the paper [V. Pragadeeswarar, M. Marudai, Best proximity points: approximation and optimization in partially ordered metric spaces, *Optim. Lett.* 7 (2013), 1883–1892] the results are not real generalizations but particular cases of existing fixed point theorems in the literature.

MSC(2010): 90C26, 90C30.

Keywords: Partially ordered set; Proximally increasing mapping; Ordered proximal contraction; Best proximity point.

1. Introduction

Let (X, \preceq) be a partially ordered set. A self mapping $T : X \rightarrow X$ is said to be *monotone nondecreasing* if $T(x) \preceq T(y)$ whenever $x, y \in X, x \preceq y$. In 2005 the following fixed point theorem was established by Nieto and Rodri'guez-Lo'pez for monotone nondecreasing mappings which can be considered as an extension of the *Banach contraction principle*. We will provide a brief proof here since the main ideas will be used in the sequel.

Theorem 1.1. ([1]) *Let (X, \preceq) be a partially ordered set and $T : X \rightarrow X$ be a self mapping which is monotone nondecreasing. Assume that there is a metric d on X such that (X, d) is a complete metric space and X satisfies the condition*

(1.1) *if a nondecreasing sequence $\{x_n\} \rightarrow x \in X$, then $x_n \preceq x, \forall n$.*

Suppose that there exists $\alpha \in [0, 1[$ such that $d(Tx, Ty) \leq \alpha d(x, y)$ for every $x, y \in X$ with $x \preceq y$. If there exists $x_0 \in X$ with $x_0 \preceq T(x_0)$, then T has

Date: Received: February 11, 2021, Accepted: May 29, 2021.

*Corresponding author.

a fixed point. Moreover, if we define $x_n = Tx_{n-1}$ for all $n \in \mathbb{N}$, then the sequence $\{x_n\}$ converges to a fixed point of T .

Proof. Since $x_0 \in X$ with $x_0 \preceq T(x_0)$ and T is monotone nondecreasing, the Picard's iteration sequence $\{T^n(x_0)\}$ is increasing. It now follows from the assumption on the mapping T that there exists $\alpha \in [0, 1[$ such that

$$d(T^{n+1}x_0, T^n x_0) \leq \alpha d(T^n x_0, T^{n-1}x_0), \quad \forall n \in \mathbb{N},$$

that is, $\{T^n(x_0)\}$ is a Cauchy sequence and so converges to an element $p \in X$. By using (1) we conclude that $x_n \preceq p$ for all $n \in \mathbb{N}$. We now have

$$d(T^{n+1}x_0, Tp) \leq \alpha d(T^n x_0, p) \xrightarrow{(n \rightarrow \infty)} 0,$$

which ensures that p is a fixed point of T . \square

Throughout this article we denote by Ψ the class of the *altering distance functions* $\psi : [0, \infty) \rightarrow [0, \infty)$ which satisfy the following conditions:

- (i) ψ is continuous and nondecreasing;
- (ii) $\psi(t) = 0$ if and only if $t = 0$.

This class of functions was first introduced in [6].

In [5] Harjani and Sadarangani established the following extension of Theorem 1.1 by using altering distance functions as control functions on contractive conditions.

Theorem 1.2. ([5]) *Let (X, \preceq) be a partially ordered set and suppose that there exists a metric d in X such that (X, d) is a complete metric space and X satisfies the condition (1) of Theorem 1.1. Let $T : X \rightarrow X$ be a monotone nondecreasing self mapping such that*

$$(1.2) \quad \psi(d(Tx, Ty)) \leq \psi(d(x, y)) - \varphi(d(x, y)), \quad \forall x, y \in X \text{ with } x \preceq y,$$

where $\psi, \varphi \in \Psi$. If there exists $x_0 \in X$ with $x_0 \preceq T(x_0)$, then T has a fixed point. Moreover, if we define $x_n = Tx_{n-1}$ for all $n \in \mathbb{N}$, then the sequence $\{x_n\}$ converges to the fixed point of T .

Recently, Theorem 1.1 and Theorem 1.2 was generalized in [9] and [7] in order to resolve an optimization problem in the setting of a metric space that is endowed with a partial order.

In this article we show that the results of [7, 9] not only are not real extensions of Theorem 1.1, Theorem 1.2 but also they are consequences of Theorem 1.1 and Theorem 1.2, respectively. We refer to [3, 4] for more related subject.

2. Preliminaries

Let (X, d) be a metric space equipped with a partial order relation " \preceq " and (A, B) be a pair of nonempty subsets of X . We use the following notions and notations in the sequel:

$$\begin{aligned} \text{dist}(A, B) &:= \inf\{d(x, y) : (x, y) \in A \times B\}, \\ A_0 &:= \{x \in A : d(x, y) = \text{dist}(A, B), \text{ for some } y \in B\}, \end{aligned}$$

$$B_0 := \{y \in B : d(x, y) = \text{dist}(A, B), \text{ for some } x \in A\},$$

We mention that a point $x^* \in A$ is said to be a *best proximity point* for a non-self mapping $T : A \rightarrow B$ provided that

$$d(x^*, Tx^*) = \text{dist}(A, B).$$

It is remarkable to note that if $x^* \in A$ is a best proximity point for the non-self mapping T , then it is a solution of the following minimization problem: Find

$$(2.1) \quad \min_{x \in A} d(x, Tx).$$

Definition 2.1. ([10]) The pair (A, B) is said to have P-property if and only if

$$\begin{cases} d(x_1, y_1) = \text{dist}(A, B), \\ d(x_2, y_2) = \text{dist}(A, B), \end{cases} \implies d(x_1, x_2) = d(y_1, y_2),$$

where $x_1, x_2 \in A_0$ and $y_1, y_2 \in B_0$.

Definition 2.2. ([8]) A non-self mapping $T : A \rightarrow B$ is said to be proximally increasing if it satisfies the condition that

$$\begin{cases} x_1 \preceq x_2, \\ d(u_1, Tx_1) = \text{dist}(A, B), \\ d(u_2, Tx_2) = \text{dist}(A, B), \end{cases} \implies u_1 \preceq u_2,$$

for all $x_1, x_2, u_1, u_2 \in A$.

Definition 2.3. ([8]) A non-self mapping $T : A \rightarrow B$ is said to be an ordered proximal contraction if there exists a non-negative real number $\alpha < 1$ such that

$$\begin{cases} x_1 \preceq x_2, \\ d(u_1, Tx_1) = \text{dist}(A, B), \\ d(u_2, Tx_2) = \text{dist}(A, B), \end{cases} \implies d(u_1, u_2) \leq \alpha d(x_1, x_2),$$

for all $x_1, x_2, u_1, u_2 \in A$.

Definition 2.4. ([9]) Given non-self mappings $S, T : A \rightarrow B$ the pair $(S; T)$ is said to be proximally increasing if

$$\begin{cases} x \preceq y, \\ d(u, Sx) = \text{dist}(A, B), \\ d(v, Ty) = \text{dist}(A, B), \end{cases} \implies u \preceq v,$$

for all $x, u \in A, y, v \in B$.

Definition 2.5. ([9]) Given non-self mappings $S, T : A \rightarrow B$ the pair $(S; T)$ is form an ordered proximal cyclic contraction if there exists a non-negative

real number $\beta < 1$ such that

$$\begin{cases} x \preceq y, \\ d(u, Sx) = \text{dist}(A, B), \\ d(v, Ty) = \text{dist}(A, B), \end{cases} \implies d(u, v) \leq \beta d(x, y) + (1 - \beta) \text{dist}(A, B),$$

for all $x, u \in A, y, v \in B$.

Here we state the main results of [7, 9].

Theorem 2.6. (see Theorem 3.1 of [9]) *Let X be a nonempty set such that (X, \preceq) is a partially ordered set and (X, d) is a complete metric space. Let A and B be non-void closed subsets of the metric space (X, d) such that A_0 is nonempty. Let $S, T : A \rightarrow B$ and $g : A \cup B \rightarrow A \cup B$ satisfy the following conditions:*

- (i) S and T are proximally increasing, ordered proximal contractions;
- (ii) $S(A_0) \subseteq B_0$ and $T(B_0) \subseteq A_0$;
- (iii) g is a surjective isometry, its inverse is an increasing mapping, $A_0 \subseteq g(A_0)$ and $B_0 \subseteq g(B_0)$;
- (iv) The pair $(S; T)$ forms a proximally increasing, ordered proximal cyclic contraction.
- (v) There exist elements $x_0, x_1 \in A_0$ and $y_0, y_1 \in B_0$ such that

$$d(gx_1, Sx_0) = \text{dist}(A, B) = d(gy_1, Ty_0),$$

where $x_0 \preceq x_1, y_0 \preceq y_1$ and $x_0 \preceq y_0$;

- (vi) The sets A and B satisfy the condition (1) of Theorem 1.1.

Then there exists an element $(x^*, y^*) \in A \times B$ such that

$$d(gx^*, Sx^*) = d(gy^*, Ty^*) = d(x^*, y^*) = \text{dist}(A, B).$$

Further the sequence $(\{x_n\}, \{y_n\})$ in $A_0 \times B_0$ defined by

$$d(gx_{n+1}, Sx_n) = \text{dist}(A, B) = d(gy_{n+1}, Ty_n), \quad \forall n \in \mathbb{N} \cup \{0\},$$

converges to the element (x^*, y^*) .

Theorem 2.7. (see Theorems 2.1 and 2.2 of [7]) *Let X be a nonempty set such that (X, \preceq) is a partially ordered set and (X, d) is a complete metric space. Let A and B be non-void closed subsets of the metric space (X, d) such that A_0 is nonempty. Let $T : A \rightarrow B$ satisfy the following conditions:*

- (i) T is a proximally increasing such that $T(A_0) \subseteq B_0$ and (A, B) satisfies the P -property;
- (ii) there exist elements x_0 and x_1 in A_0 such that

$$x_0 \preceq x_1, \quad d(x_1, Tx_0) = \text{dist}(A, B),$$

- (iii) for all $x, y \in A$ with $x \preceq y$,

$$(2.2) \quad \psi(d(Tx, Ty)) \leq \psi(d(x, y)) - \varphi(d(x, y)),$$

where $\varphi, \psi \in \Psi$;

(iv) The set A satisfies the condition (1) of Theorem 1.1.

Then T has a best proximity point. Further the sequence $\{x_n\}$ defined by

$$d(x_{n+1}, Tx_n) = \text{dist}(A, B), \quad \forall n \in \mathbb{N} \cup \{0\},$$

converges to the best proximity point of T .

3. Main results

Theorem 3.1. Theorem 2.6 is a straightforward consequence of Theorem 1.1.

Proof. Let $x \in A_0$. Since $Sx \in B_0$, there exists an element $u \in A_0$ such that $d(u, Sx) = \text{dist}(A, B)$. By the fact that $A_0 \subseteq g(A_0)$, we can find an element $\hat{u} \in A_0$ for which $u = g\hat{u}$ and so $d(g\hat{u}, Sx) = \text{dist}(A, B)$. It is worth noticing that if there exists another element $\check{u} \in A_0$ for which $d(g\check{u}, Sx) = \text{dist}(A, B)$, then by this reality that S is an ordered proximal contraction and g is an isometry, we obtain

$$d(\hat{u}, \check{u}) = d(g\hat{u}, g\check{u}) \leq \alpha d(x, x) = 0,$$

which implies that $\hat{u} = \check{u}$. Thus we can define a self mapping $\Pi_1 : A_0 \rightarrow A_0$ such that $d(g\Pi_1x, Sx) = \text{dist}(A, B)$ for all $x \in A_0$. By a similar argument we consider the self mapping $\Pi_2 : B_0 \rightarrow B_0$ for which $d(g\Pi_2y, Ty) = \text{dist}(A, B)$ for any $y \in B_0$. We have the following observations about the mappings Π_i for $i \in \{1, 2\}$.

♣ Let $x_1, x_2 \in A_0$ be such that $x_1 \preceq x_2$. Then

$$\begin{cases} d(g\Pi_1x_1, Sx_1) = \text{dist}(A, B), \\ d(g\Pi_1x_2, Sx_2) = \text{dist}(A, B). \end{cases}$$

Since S is a proximally increasing, $g\Pi_1x_1 \preceq g\Pi_1x_2$. Since g^{-1} is increasing, we must have $\Pi_1x_1 \preceq \Pi_1x_2$, that is, Π_1 is monotone nondecreasing. Equivalently, we can see that Π_2 is also monotone nondecreasing.

♣ Let $x_1, x_2 \in A_0$ be such that $x_1 \preceq x_2$. Then

$$\begin{cases} d(g\Pi_1x_1, Sx_1) = \text{dist}(A, B), \\ d(g\Pi_1x_2, Sx_2) = \text{dist}(A, B). \end{cases}$$

Since S is an ordered proximal contraction, there exists $\alpha \in [0, 1)$ such that

$$d(\Pi_1x_1, \Pi_1x_2) = d(g\Pi_1x_1, g\Pi_1x_2) \leq \alpha d(x_1, x_2).$$

Similarly, if $y_1, y_2 \in B_0$ with $y_1 \preceq y_2$, then

$$d(\Pi_2y_1, \Pi_2y_2) \leq \alpha d(y_1, y_2).$$

♣ By the assumption (v) of Theorem 2.6, there exist $x_0, x_1 \in A_0$ and $y_0, y_1 \in B_0$ with $x_0 \preceq x_1$ and $y_0 \preceq y_1$ such that $d(gx_1, Sx_0) = \text{dist}(A, B) = d(gy_1, Ty_0)$. Besides, by the definition of the mapping Π_1 , we have $d(g\Pi_1x_0, Sx_0) = \text{dist}(A, B)$. Because of the fact that S is an ordered proximal contraction, we

conclude that $x_1 = \Pi_1 x_0$ and so $x_0 \preceq \Pi_1 x_0$. Similarly, we obtain $y_0 \preceq \Pi_2 y_0$.
 ♣ Now define the mapping $\Pi : A_0 \cup B_0 \rightarrow A_0 \cup B_0$ with

$$\Pi z = \begin{cases} \Pi_1 z & \text{if } z \in A_0, \\ \Pi_2 z & \text{if } z \in B_0. \end{cases}$$

Then $\Pi(A_0) \subseteq A_0$ and $\Pi(B_0) \subseteq B_0$, that is, Π is *noncyclic* on $A_0 \cup B_0$. Let $(x, y) \in A_0 \times B_0$ be such that $x \preceq y$. Then we have

$$\begin{cases} d(g\Pi x, Sx) = \text{dist}(A, B), \\ d(g\Pi y, Ty) = \text{dist}(A, B). \end{cases}$$

Since the pair $(S; T)$ forms an ordered proximal cyclic contraction, we obtain

$$d(\Pi x, \Pi y) = d(g\Pi_1 x, g\Pi_2 y) \leq \beta d(x, y) + (1 - \beta) \text{dist}(A, B).$$

♣ For the considered elements $(x_0, y_0), (x_1, y_1) \in A_0 \times B_0$ which satisfy the condition (v) since $x_0 \preceq \Pi_1 x_0$ and Π_1 is monotone nondecreasing, the sequence $\{\Pi_1^n x_0\}$ is increasing. Similarly, the sequence $\{\Pi_2^n y_0\}$ is also increasing. It now follows from the proof of Theorem 1.1 that the sequences $\{\Pi_1^n x_0\}$ and $\{\Pi_2^n y_0\}$ are Cauchy. Let $(x^*, y^*) \in A \times B$ be such that

$$\Pi_1^n x_0 \rightarrow x^*, \quad \Pi_2^n y_0 \rightarrow y^*.$$

If we prove that $(x^*, y^*) \in A_0 \times B_0$ then by a similar argument of the proof of Theorem 1.1 we deduce that x^* and y^* are the fixed points of Π_1 and Π_2 , respectively. To show this, we note that since $x_0 \preceq y_0$ we have

$$d(\Pi x_0, \Pi y_0) \leq \beta d(x_0, y_0) + (1 - \beta) \text{dist}(A, B).$$

Since

$$\begin{cases} d(g\Pi x_0, Sx_0) = \text{dist}(A, B), \\ d(g\Pi y_0, Ty_0) = \text{dist}(A, B), \end{cases}$$

and the pair $(S; T)$ forms a proximally increasing, we conclude that $g\Pi x_0 \preceq g\Pi y_0$. By the fact that g^{-1} is increasing, $\Pi x_0 \preceq \Pi y_0$. Again, since the pair $(S; T)$ forms an ordered proximal cyclic contraction, we obtain

$$\begin{aligned} d(\Pi^2 x_0, \Pi^2 y_0) &\leq \beta d(\Pi x_0, \Pi y_0) + (1 - \beta) \text{dist}(A, B) \\ &\leq \beta^2 d(x_0, y_0) + (1 - \beta^2) \text{dist}(A, B). \end{aligned}$$

Continuing this process and by induction, we conclude that

$$d(\Pi^n x_0, \Pi^n y_0) \leq \beta^n d(x_0, y_0) + (1 - \beta^n) \text{dist}(A, B).$$

Letting $n \rightarrow \infty$ in above inequality, we obtain $d(x^*, y^*) = \text{dist}(A, B)$, that is, $(x^*, y^*) \in A_0 \times B_0$. Hence,

$$\begin{aligned} d(gx^*, Sx^*) &= d(g\Pi_1 x^*, Sx^*) = \text{dist}(A, B), \\ d(gy^*, Ty^*) &= d(g\Pi_2 y^*, Ty^*) = \text{dist}(A, B), \\ d(x^*, y^*) &= \text{dist}(A, B). \end{aligned}$$

Finally, if for each $n \in \mathbb{N}$ we set $x_n = \Pi^n x_0$ and $y_n = \Pi^n y_0$, then

$$\begin{aligned} d(gx_{n+1}, Sx_n) &= \text{dist}(A, B), \\ d(gy_{n+1}, Ty_n) &= \text{dist}(A, B), \\ (x_n, y_n) &\rightarrow (x^*, y^*). \end{aligned}$$

□

Theorem 3.2. *Theorem 2.7 is a straightforward consequence of Theorem 1.2.*

Proof. Since the pair (A, B) has the P-property, it follows from Lemma 3.1 of [2] that both A_0 and B_0 are closed. Moreover, if $x \in A_0$, then there exists an element $v \in B_0$ such that $d(x, v) = \text{dist}(A, B)$. We note that if there is another element $v' \in B_0$ for which $d(x, v') = \text{dist}(A, B)$, then from the fact that (A, B) has the P-property, we must have $v = v'$. So, we can define a mapping $g : A_0 \rightarrow B_0$ such that

$$d(x, gx) = \text{dist}(A, B), \quad \forall x \in A_0.$$

It is worth noticing that for any $u_1, u_2 \in A_0$, we have $d(u_1, gu_1) = \text{dist}(A, B) = d(u_2, gu_2)$ which ensures that

$$d(u_1, u_2) = d(gu_1, gu_2), \quad \forall u_1, u_2 \in A_0,$$

that is, g is an isometry. Hence, g is a bijective isometry mapping. Now consider the self-mapping $g^{-1}T : A_0 \rightarrow A_0$. Here, we check the conditions of Theorem 1.1 for the self mapping $g^{-1}T : A_0 \rightarrow A_0$.

♠ Let $x, y \in A_0$ be such that $x \preceq y$. Since g^{-1} is an isometry, we conclude that

$$\psi\left(d((g^{-1}T)x, (g^{-1}T)y)\right) = \psi(d(Tx, Ty)) \leq \psi(d(x, y)) - \varphi(d(x, y)),$$

where $\varphi, \psi \in \Psi$.

♠ It follows from the assumption (ii) of Theorem 2.7 that there exist the elements $x_0, x_1 \in A_0$ such that $x_0 \preceq x_1$ and $d(x_1, Tx_0) = \text{dist}(A, B)$. By the fact that $d(x_1, gx_1) = \text{dist}(A, B)$ and that (A, B) has the P-property, we obtain $gx_1 = Tx_0$ and so, $x_1 = (g^{-1}T)x_0$ which implies that

$$x_0 \preceq (g^{-1}T)x_0.$$

♠ Let $x, y \in A_0$ be such that $x \preceq y$. Since $T(A_0) \subseteq B_0$ there are two points $u, v \in A_0$ such that

$$d(u, Tx) = \text{dist}(A, B) = d(v, Ty).$$

Because T is proximally increasing, we must have $u \preceq v$. Besides, from the definition of the mapping g we have $gu = Tx$ and $gv = Ty$ and hence

$$(g^{-1}T)x = u \preceq v = (g^{-1}T)y,$$

which implies that the self mapping $g^{-1}T$ is monotone nondecreasing.

Thereby, all of the assumptions of Theorem 1.1 hold and the self mapping

$g^{-1}T : A_0 \rightarrow A_0$ has a fixed point, called $x^* \in A_0$, that is, $g^{-1}Tx^* = x^*$ which ensures that $Tx^* = gx^*$. Hence,

$$d(x^*, Tx^*) = d(x^*, gx^*) = \text{dist}(A, B).$$

On the other hand if we define $x_n = (g^{-1}T)x_{n-1}$ for any $n \in \mathbb{N}$, then $x_n \rightarrow x^*$. In this case we have $gx_n = Tx_{n-1}$ and so

$$d(x_n, Tx_{n-1}) = d(x_n, gx_n) = \text{dist}(A, B),$$

and the result follows. \square

4. Concluding Remarks

It was proved by Sadiq Basha that in the setting of compete partially ordered metric spaces a pair of ordered proximal contractions which are proximally increasing has a common best proximity point (see Theorem 2.6). Moreover, an existence and convergence result of a best proximity point for proximally increasing nonself mappings was established by Pragadeeswarar and Maruda using a geometric concept of P-property (see Theorem 2.7).

We have proved that these existence results are straightforward consequences of Theorem 1.1 and Theorem 1.2, respectively.

REFERENCES

- [1] Nieto, J.J., Rodri'guez-Lo'pez, R., *Contractive mapping theorems in partially ordered sets and applications to ordinary differential equations*, Order, **22** (2005), 223-239.
- [2] Gabeleh, M.: *Proximal weakly contractive and proximal nonexpansive non-self-mappings in metric and Banach spaces*, J. Optim. Theory Appl., **158** (2013), 615-625.
- [3] Gabeleh, M., Markin, J.: *A note on the paper "Best proximity point results for p-proximal contractions*, Acta Math. Hungar., (2021), doi.org/10.1007/s10474-021-01139-5.
- [4] Gabeleh, M., Vetro, C.: *A note on best proximity point theory using proximal contractions*, J. Fixed Point Theory Appl., **20** (2018), doi.org/10.1007/s11784-018-0624-4.
- [5] Harjani, J., Sadarangani, K.: *Generalized contractions in partially ordered metric spaces and applications to ordinary differential equations*, Nonlinear Anal., **72** (2010), 1188-1197.
- [6] Khan, M.S., Swaleh, M., Sessa, S.: *Fixed point theorems by altering distances between the points*, Bull. Austral. Math. Soc., **30** (1984), 1-9.
- [7] Pragadeeswarar, V., Marudai, M.: *Best proximity points: approximation and optimization in partially ordered metric spaces*, Optim. Lett., **7** (2013), 1883-1892.
- [8] Sadiq Basha, S., *Discrete optimization in partially ordered sets*, J. Global Optim., **54** (2012), 511-517.
- [9] Sadiq Basha, S., *Global optimization in metric spaces with partial orders*, Optimization, **63** (2014), 817-825.
- [10] Sankar Raj, V., *A best proximity point theorem for weakly contractive non-self-mappings*, Nonlinear Anal., **74** (2011), 4804-4808.

(Moosa Gabeleh) DEPARTMENT OF MATHEMATICS, AYATOLLAH BOROUJERDI UNIVERSITY, BOROUJERD, IRAN.

Email address: Gabeleh@abru.ac.ir

(Jack Markin) 1440 8TH ST. GOLDEN, CO 80401, USA; JMARKIN@NEWMEXICO.COM

Email address: jmarkin@newmexico.com